

MEIAMUT, D.L., kand.tekhn.nauk

Construction of hydraulic-fill dams in land reclamation projects.
Trudy VNIIGIM 32:175-182 '59. (MIRA 13:8)
(Dams)

MELAMUT, D.L., kand.tekhn.nauk

Using hydraulic fill methods in constructing a dam on the Murgab
River. Transp.stroi. 10 no.6:29-32 Je '60. (MIRA 13:7)
(Murgab River--Dams)

MELAMUT, D.L., kand.tekhn.nauk; YEROKHIN, V.D., inzh.

Using hydraulic fill methods on one side only in building the dam
of a reservoir. Transp. stroi. ll no.1:18-20 Ja '61.

(MIRA 14:1)

(Dams)

MELAMUT, D.L., kand.tekhn.nauk; SILAGADZE, V.A., inzh.

Cofferdamming by breaking up the drop. Gidr. i mel. 14 no.8:
25-30 Ag '62. (MIRA 15:9)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut gidrotekhniki
i melioratsii.

(Cofferdams)

MELAMUT, D.L., kand.tekhn.nauk; NIKOLAYEV, V.M., insh.

Method for calculating the stability of slopes for narrow-profile
hydraulic-fill dams. Gidr. i mel. 14 no.12:43-50 D '62.

(MIRA 16:5)

(Dams)

MEIA'UT, David Lazarevich, kand. tekhn. nauk; NIKOLAYEV, Vasilii
Mikhaylovich, kand. tekhn. nauk; TSAREVSKIY, A.M., retsenzent,
AFANAS'YEV, B.P., red. izd-va; RODIONOVA, V.M., tekhn. red.

[Hydraulic filling of narrow-profile dams and small dams in
agricultural construction work] Namyv uzkoprofil'nykh damb i
malykh plotin v sel'skom stroitel'stve. Moskva, Gosstroizdat,
1963. 241 p. (MIRA 16:7)

1. Vsesoyuznaya akademiya sel'skokhozyaystvennykh nauk imeni
V.I. Lenina (for TSarevskiy).

(Dams)

MELAMUT, D.L., kand. tekhn. nauk; SILAGADZE, V.A., inzh.

Cofferdamming channels with local soil. Trans. stroi. 13
no.8:36-39 Ag '63. (MIRA 17:2)

MELAMUT, D.L., kand. tekhn. nauk; MENTYUKOV, V.P., inzh.

Hydraulic-file construction of dams from gravelly soils
with formation of an antiseepage core. Gidr. i mel. 15
no.8:26-32 Ag '63. (MIRA 16:8)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut gidro-
tekhniki i melioratsii im. Kostyakova.

KALIZHNYUK, S.K., inzh.; MELAMUT, D.L., kand.tekhn.nauk; SILAGADZE, V.A.,
inzh.

Results of cofferdamming with sand and gravel. Gidr.stroi. 34
no.11:10-15 N '63. (MIRA 17:3)

KOSTYAKOVA, A.N., nauchn. sotr.; MELAMUT, D.L., kand. tekhn.
nauk, nauchn. sotr.; KENTYUKOV, V.P., inzh., nauchn.
sotr.

[Hydraulic fill of dams composed of gravelly soil] Namyv
plotin iz gravelistyykh gruntov. Moskva, Sel'khozizdat,
1963. 12 p. (MIRA 17:9)

1. Russia (1923- U.S.S.R.) Ministerstvo sel'skogo kho-
zyaystva. Upravleniye nauki, propagandy i vnedreniya pe-
redovogo opyta. 2. Laboratoriya gidromekhanizatsii zem-
lyanykh rabot Vsesoyuznogo nauchno-issledovatel'skogo
instituta gidrotekhniki i melioratsii.

MULLER, R.A., kand.tekhn.nauk; YUSHIN, A.I., inzh.; MELAMUT, L.Sh., inzh.

Temporary technical specifications for planning and constructing buildings and structures on ground located over mines. Shakht. stroi.

4 no. 5:29-30 My '60.

(Foundations) (Soil mechanics)

(MIRA 14:4)

KOLBENKOV, S.P.; MEDYANTSEV, A.N.; IOFIS, M.A.; KOROTKOV, M.V.;
MULLER, R.A.; YUSHIN, A.I.; MELAMUT, L.Sh.; KARGIN, G.P.;
GERTNER, P.F.; ZARETSKIY, K.S.; CHECHKOV, L.V., red.izd-
va; MAKSIMOVA, V.V., tekhn. red.

[Designing, constructing, and protecting buildings and
structures on foundations undercut by mining] Proektiro-
vanie, stroitel'stvo i okhrana zdaniy i sooruzheniy na pod-
rabatyvaemykh territoriyakh. Moskva, Gosgortekhnizdat,
1963. 451 p. (MIRA 16:8)

(Earth movements and building)

1ST AND 2ND ORDERS

PROCESSES AND PROPERTIES INDEX

3RD AND 4TH ORDERS

MELAN, H
F

522. NEW TENDENCIES IN DEVELOPMENT OF CONSTRUCTION OF GAS AND STEAM TURBINES WITH SPECIAL REFERENCE TO AUSTRIAN POWER STATION PROJECTS. Melan, H. (Öst. Z. Elektrizitätswirtsch., Dec. 1949, vol. 2, 292-301). A review with 17 diagrams and illustrations.

SA

ASME-SEA METALLURGICAL LITERATURE CLASSIFICATION

REGIONAL INDEX

1949-1950

1951-1952

1953-1954

1955-1956

1957-1958

1959-1960

1961-1962

1963-1964

1965-1966

1967-1968

1969-1970

1971-1972

1973-1974

1975-1976

1977-1978

1979-1980

1981-1982

1983-1984

1985-1986

1987-1988

1989-1990

1991-1992

1993-1994

1995-1996

1997-1998

1999-2000

2001-2002

2003-2004

2005-2006

2007-2008

2009-2010

2011-2012

2013-2014

2015-2016

2017-2018

2019-2020

2021-2022

2023-2024

2025-2026

2027-2028

2029-2030

2031-2032

2033-2034

2035-2036

2037-2038

2039-2040

2041-2042

2043-2044

2045-2046

2047-2048

2049-2050

2051-2052

2053-2054

2055-2056

2057-2058

2059-2060

2061-2062

2063-2064

2065-2066

2067-2068

2069-2070

2071-2072

2073-2074

2075-2076

2077-2078

2079-2080

2081-2082

2083-2084

2085-2086

2087-2088

2089-2090

2091-2092

2093-2094

2095-2096

2097-2098

2099-2100

2101-2102

2103-2104

2105-2106

2107-2108

2109-2110

2111-2112

2113-2114

2115-2116

2117-2118

2119-2120

2121-2122

2123-2124

2125-2126

2127-2128

2129-2130

2131-2132

2133-2134

2135-2136

2137-2138

2139-2140

2141-2142

2143-2144

2145-2146

2147-2148

2149-2150

2151-2152

2153-2154

2155-2156

2157-2158

2159-2160

2161-2162

2163-2164

2165-2166

2167-2168

2169-2170

2171-2172

2173-2174

2175-2176

2177-2178

2179-2180

2181-2182

2183-2184

2185-2186

2187-2188

2189-2190

2191-2192

2193-2194

2195-2196

2197-2198

2199-2200

2201-2202

2203-2204

2205-2206

2207-2208

2209-2210

2211-2212

2213-2214

2215-2216

2217-2218

2219-2220

2221-2222

2223-2224

2225-2226

2227-2228

2229-2230

2231-2232

2233-2234

2235-2236

2237-2238

2239-2240

2241-2242

2243-2244

2245-2246

2247-2248

2249-2250

2251-2252

2253-2254

2255-2256

2257-2258

2259-2260

2261-2262

2263-2264

2265-2266

2267-2268

2269-2270

2271-2272

2273-2274

2275-2276

2277-2278

2279-2280

2281-2282

2283-2284

2285-2286

2287-2288

2289-2290

2291-2292

2293-2294

2295-2296

2297-2298

2299-2300

2301-2302

2303-2304

2305-2306

2307-2308

2309-2310

2311-2312

2313-2314

2315-2316

2317-2318

2319-2320

2321-2322

2323-2324

2325-2326

2327-2328

2329-2330

2331-2332

2333-2334

2335-2336

2337-2338

2339-2340

2341-2342

2343-2344

2345-2346

2347-2348

2349-2350

2351-2352

2353-2354

2355-2356

2357-2358

2359-2360

2361-2362

2363-2364

2365-2366

2367-2368

2369-2370

2371-2372

2373-2374

2375-2376

2377-2378

2379-2380

2381-2382

2383-2384

2385-2386

2387-2388

2389-2390

2391-2392

2393-2394

2395-2396

2397-2398

2399-2400

2401-2402

2403-2404

2405-2406

2407-2408

2409-2410

2411-2412

2413-2414

2415-2416

2417-2418

2419-2420

2421-2422

2423-2424

2425-2426

2427-2428

2429-2430

2431-2432

2433-2434

2435-2436

2437-2438

2439-2440

2441-2442

2443-2444

2445-2446

2447-2448

2449-2450

2451-2452

2453-2454

2455-2456

2457-2458

2459-2460

2461-2462

2463-2464

2465-2466

2467-2468

2469-2470

2471-2472

2473-2474

2475-2476

2477-2478

2479-2480

2481-2482

2483-2484

2485-2486

2487-2488

2489-2490

2491-2492

2493-2494

2495-2496

2497-2498

2499-2500

2501-2502

2503-2504

2505-2506

2507-2508

2509-2510

2511-2512

2513-2514

2515-2516

2517-2518

2519-2520

2521-2522

2523-2524

2525-2526

2527-2528

2529-2530

2531-2532

2533-2534

2535-2536

2537-2538

2539-2540

2541-2542

2543-2544

2545-2546

2547-2548

2549-2550

2551-2552

2553-2554

2555-2556

2557-2558

2559-2560

2561-2562

2563-2564

2565-2566

2567-2568

2569-2570

2571-2572

2573-2574

2575-2576

2577-2578

2579-2580

2581-2582

2583-2584

2585-2586

2587-2588

2589-2590

2591-2592

2593-2594

2595-2596

2597-2598

2599-2600

2601-2602

2603-2604

2605-2606

2607-2608

2609-2610

2611-2612

2613-2614

2615-2616

2617-2618

2619-2620

2621-2622

2623-2624

2625-2626

2627-2628

2629-2630

2631-2632

2633-2634

2635-2636

2637-2638

2639-2640

2641-2642

2643-2644

2645-2646

2647-2648

2649-2650

2651-2652

2653-2654

2655-2656

2657-2658

2659-2660

2661-2662

2663-2664

2665-2666

2667-2668

2669-2670

2671-2672

2673-2674

2675-2676

2677-2678

2679-2680

2681-2682

2683-2684

2685-2686

2687-2688

2689-2690

2691-2692

2693-2694

2695-2696

2697-2698

2699-2700

2701-2702

2703-2704

2705-2706

2707-2708

2709-2710

2711-2712

2713-2714

2715-2716

2717-2718

2719-2720

2721-2722

2723-2724

2725-2726

2727-2728

2729-2730

2731-2732

2733-2734

2735-2736

2737-2738

2739-2740

2741-2742

2743-2744

2745-2746

2747-2748

2749-2750

2751-2752

2753-2754

2755-2756

2757-2758

2759-2760

2761-2762

2763-2764

2765-2766

2767-2768

2769-2770

2771-2772

2773-2774

2775-2776

2777-2778

2779-2780

2781-2782

2783-2784

2785-2786

2787-2788

2789-2790

2791-2792

2793-2794

2795-2796

2797-2798

2799-2800

2801-2802

2803-2804

2805-2806

2807-2808

2809-2810

2811-2812

2813-2814

2815-2816

2817-2818

2819-2820

2821-2822

2823-2824

2825-2826

2827-2828

2829-2830

2831-2832

2833-2834

2835-2836

2837-2838

2839-2840

2841-2842

2843-2844

2845-2846

2847-2848

2849-2850

2851-2852

2853-2854

2855-2856

2857-2858

2859-2860

2861-2862

2863-2864

2865-2866

2867-2868

2869-2870

2871-2872

2873-2874

2875-2876

2877-2878

2879-2880

2881-2882

2883-2884

2885-2886

2887-2888

2889-2890

2891-2892

2893-2894

2895-2896

2897-2898

2899-2900

2901-2902

2903-2904

2905-2906

2907-2908

2909-2910

2911-2912

2913-2914

2915-2916

2917-2918

2919-2920

2921-2922

2923-2924

2925-2926

2927-2928

2929-2930

2931-2932

2933-2934

2935-2936

2937-2938

2939-2940

2941-2942

2943-2944

2945-2946

2947-2948

2949-2950

2951-2952

2953-2954

2955-2956

2957-2958

2959-2960

2961-2962

2963-2964

2965-2966

2967-2968

2969-2970

2971-2972

2973-2974

2975-2976

2977-2978

2979-2980

2981-2982

2983-2984

2985-2986

2987-2988

2989-2990

2991-2992

2993-2994

2995-2996

2997-2998

2999-3000

MELAN, H.

"Propositions for the thermodynamics of nuclear power plants."

p. 517 (Energia Es Atomtechnika) Vol. 10, no. 8/10, Dec. 1957
Budapest, Hungary

SO: Monthly Index of East European Accessions (EJAI) LC. Vol. 7, no. 4,
April 1958

MELAN, K.G., prof. (Rostok, Germanskaya Demokraticheskaya Respublika)

In the name of peace and progress. Zdorov'e 5 no.9:9
S '59. (MIRA 12:11)
(GERMANY, EAST--PUBLIC HEALTH)

S/081/62/000/023/100/120
B101/B186

AUTHOR: Meland, Thor

TITLE: Friction and wear of plastics

PERIODICAL: Referativnyy zhurnal. Khimiya, no. 23, 1962, 693, abstract
23P199 (Tekn. ukebl., v. 109, no. 14, 1962, 351 - 360 [Norweg.;
summary in Eng.])

TEXT: Friction (F) and wear of plastics have been studied as functions of their macromolecular structure. Relations were established between F of dry gliding, the adhesion properties and the viscosity of the plastics (particularly the degree of crystallinity); also between the coefficient of F and the gliding velocity, the specific pressure, temperature, and braking time. The function of rolling F versus velocity, and the factor of dynamic losses versus frequency, were clarified. Characteristics for choosing wear-resistant plastics were determined. [Abstracter's note: Complete translation.]

Card 1/1

MELANEK, I.

"What Our Technology Gave to the World; A Comment on the Exhibit of the National Museum of Technology in Prague." p.315

"Appraisal of the Book Design and Organization of Machine Shops." Trans. from the Russian. p. 318 (Strojirenstvl, Vol. 3, no. 4, Apr. 1953, Praha)

SO: Monthly List of ^{East European} ~~Russian~~ Accessions, Vol. 3, No. 3, Library of Congress, March 1954, Uncl.

MELANICH, N. (Moskva)

In search for innovations. Izobr.i rats. no.12:34 D '60.

(MIRA 13:12)

(Moscow--Technological innovations)

MELANIFIDI, G.F.

DT-1 and IT-2 thermostat. M.M. 1963. 3-3.

'63.

(MIR 17:8)

1. Gruzinskiy filial Vnesheconombank
i proyektno-konstruktor. op. r. 1963. 1-1.
neftyanoy i gazovoy prom. 1963.

MELANIFIDI, G.F.

Automatic gas spray on a jet with automatic ignition. Gaz. delo
no.10:47-49 '63. (MIRA 13:4)

1. Cktyabr'skiy filial Vsesoyuznogo nauchno-issledovatel'skogo i
proyektiro-konstruktorskogo instituta kompleksnoy avtomatizatsii
neftyanoy i gazovoy promyshlennosti.

MELANIFIDI, G.F.

Automatic flushing of clarifying agents. Nefteprom. delo no.9:
24-26 '63. (MIRA 17:4)

1. Oktyabr'skiy filial Vsesoyuznogo nauchno-issledovatel'skogo
i projektno-konstruktorskogo instituta kompleksnoy avtomatizatsii
neftyanoy i gazovoy promyshlennosti.

MELANIFIDI, G.F.

Automatic control of the water level in blenders. Mash. i neft.
obor. no.10:18-20 '63. (MIRA 17:4)

1. Filial Vsesoyuznogo nauchno-issledovatel'skogo i proyektno-konstruktorskogo instituta kompleksnoy avtomatizatsii neftyanoy i gazovoy promyshlennosti, g. Oktyabr'skiy.

RYAPOSOV, V.V.; MELANIFIDI, G.F.

Stop device for compressors with gas motors. Gaz. delo no.10:
49-50 '63. (MIRA 17:4)

1. Oktyabr'skiy filial Vsesoyuznogo nauchno-issledovatel'skogo i
proyektno-konstruktorskogo instituta kompleksnoy avtomatizatsii
neftyanoy i gazovoy promyshlennosti.

MELANIFIDI, G.F.

Introducing automatic control into the reagent industry.
Nefteprom. delc no.4:21-23 '64. (MIRA 17:6)

1. Oktyabr'skiy filial Vsesoyuznogo nauchno-issledovatel'-
skogo i proyektno-konstruktorskogo instituta kompleksnoy
avtomatizatsii neftyanoy i gazovoy promyshlennosti.

MELANIFIDI, G.F.

DU-3 leak detector. Mash. i neft. obor. no.8:30 '64. (MIRA 17:11)

1. Oktyabr'skiy filial Vsesoyuznyy nauchno-issledovatel'skiy i proyekt-no-konstruktorskiy institut kompleksnoy avtomatizatsii neftyanoy i gazovoy promyshlennosti.

MELANIFIDI, G.F.

Automatic coagulant proportioning. Mash. i nef. sbor. no. 2:
32-33 '64. (MIRA 17:11)

1. Oktyabr'skiy filial Vsesoyuznogo nauchno-issledovatel'skogo i
proyektno-konstruktorskogo instituta kompleksnoy avtomatizatsii
neftyanoy i gazovoy promyshlennosti.

MELANIFIDI, G.F.

Automatic control of the fast filters of a water purification station. Mash. i neft. obor. no.10#28-30 '64 (MIRA 18:1)

1. Otktyabr'skiy filial Vsesoyuznogo nauchno-issledovatel'skogo i proyektno-konstruktorskogo instituta kompleksnoy avtomatizatsii neftyanoy i gazovoy promyshlennosti.

MELANIFIDI, G.F.

Automatiŝ control of the dynamic level in siphon water-supply
wells. Nefteprom. delo no.4:30-31 '65. (MIRA 18:6)

1. Okt'yabr'skiy filial Vsesoyuznogo nauchno-issledovatel'skogo i
proyektno-konstrukterskogo instituta kompleksnoy avtomatizatsii
neftyanoy i gazovoy promyshlennosti.

BC

A-2

Investigation of materials in the laboratory of the U.S. Army, Corps of Engineers, Fort Belvoir, Illinois, and the examination of a series of materials are reported by H. J. E. [illegible]

ASB-66A METALLURGICAL LITERATURE CLASSIFICATION

FORM 100-100

FORM 100-100

Ca

PROCESSING AND PROPERTIES INDEX

The coloring of magnesium-iron micas. N. M. Melankholin. *Trav. lab. crist. acad. sci. U.R.S.S.* 1940, No. 2, 61-72; *Mineralog. Abstracts* 9, 197 (1946). Absorption curves of several phlogopites and biotites, compared with those of FeSO_4 , epidote, and carnallite, point to Fe_2O_3 as the coloring matter. The influence of FeO is shown only in the red region of the spectrum. Variations shown by some phlogopites are attributed to Ti. M. F.

ASME METALLURGICAL LITERATURE CLASSIFICATION

SECTION ONE ONE

SECTION TWO TWO

SECTION THREE THREE

SECTION FOUR FOUR

SECTION FIVE FIVE

SECTION SIX SIX

SECTION SEVEN SEVEN

SECTION EIGHT EIGHT

SECTION NINE NINE

SECTION TEN TEN

SECTION ELEVEN ELEVEN

SECTION TWELVE TWELVE

SECTION THIRTEEN THIRTEEN

SECTION FOURTEEN FOURTEEN

SECTION FIFTEEN FIFTEEN

SECTION SIXTEEN SIXTEEN

SECTION SEVENTEEN SEVENTEEN

SECTION EIGHTEEN EIGHTEEN

SECTION NINETEEN NINETEEN

SECTION TWENTY TWENTY

SECTION TWENTY-ONE TWENTY-ONE

SECTION TWENTY-TWO TWENTY-TWO

SECTION TWENTY-THREE TWENTY-THREE

SECTION TWENTY-FOUR TWENTY-FOUR

SECTION TWENTY-FIVE TWENTY-FIVE

SECTION TWENTY-SIX TWENTY-SIX

SECTION TWENTY-SEVEN TWENTY-SEVEN

SECTION TWENTY-EIGHT TWENTY-EIGHT

SECTION TWENTY-NINE TWENTY-NINE

SECTION THIRTY THIRTY

SECTION THIRTY-ONE THIRTY-ONE

SECTION THIRTY-TWO THIRTY-TWO

SECTION THIRTY-THREE THIRTY-THREE

SECTION THIRTY-FOUR THIRTY-FOUR

SECTION THIRTY-FIVE THIRTY-FIVE

SECTION THIRTY-SIX THIRTY-SIX

SECTION THIRTY-SEVEN THIRTY-SEVEN

SECTION THIRTY-EIGHT THIRTY-EIGHT

SECTION THIRTY-NINE THIRTY-NINE

SECTION FORTY FORTY

SECTION FORTY-ONE FORTY-ONE

SECTION FORTY-TWO FORTY-TWO

SECTION FORTY-THREE FORTY-THREE

SECTION FORTY-FOUR FORTY-FOUR

SECTION FORTY-FIVE FORTY-FIVE

SECTION FORTY-SIX FORTY-SIX

SECTION FORTY-SEVEN FORTY-SEVEN

SECTION FORTY-EIGHT FORTY-EIGHT

SECTION FORTY-NINE FORTY-NINE

SECTION FIFTY FIFTY

SECTION FIFTY-ONE FIFTY-ONE

SECTION FIFTY-TWO FIFTY-TWO

SECTION FIFTY-THREE FIFTY-THREE

SECTION FIFTY-FOUR FIFTY-FOUR

SECTION FIFTY-FIVE FIFTY-FIVE

SECTION FIFTY-SIX FIFTY-SIX

SECTION FIFTY-SEVEN FIFTY-SEVEN

SECTION FIFTY-EIGHT FIFTY-EIGHT

SECTION FIFTY-NINE FIFTY-NINE

SECTION SIXTY SIXTY

SECTION SIXTY-ONE SIXTY-ONE

SECTION SIXTY-TWO SIXTY-TWO

SECTION SIXTY-THREE SIXTY-THREE

SECTION SIXTY-FOUR SIXTY-FOUR

SECTION SIXTY-FIVE SIXTY-FIVE

SECTION SIXTY-SIX SIXTY-SIX

SECTION SIXTY-SEVEN SIXTY-SEVEN

SECTION SIXTY-EIGHT SIXTY-EIGHT

SECTION SIXTY-NINE SIXTY-NINE

SECTION SEVENTY SEVENTY

SECTION SEVENTY-ONE SEVENTY-ONE

SECTION SEVENTY-TWO SEVENTY-TWO

SECTION SEVENTY-THREE SEVENTY-THREE

SECTION SEVENTY-FOUR SEVENTY-FOUR

SECTION SEVENTY-FIVE SEVENTY-FIVE

SECTION SEVENTY-SIX SEVENTY-SIX

SECTION SEVENTY-SEVEN SEVENTY-SEVEN

SECTION SEVENTY-EIGHT SEVENTY-EIGHT

SECTION SEVENTY-NINE SEVENTY-NINE

SECTION EIGHTY EIGHTY

SECTION EIGHTY-ONE EIGHTY-ONE

SECTION EIGHTY-TWO EIGHTY-TWO

SECTION EIGHTY-THREE EIGHTY-THREE

SECTION EIGHTY-FOUR EIGHTY-FOUR

SECTION EIGHTY-FIVE EIGHTY-FIVE

SECTION EIGHTY-SIX EIGHTY-SIX

SECTION EIGHTY-SEVEN EIGHTY-SEVEN

SECTION EIGHTY-EIGHT EIGHTY-EIGHT

SECTION EIGHTY-NINE EIGHTY-NINE

SECTION NINETY NINETY

SECTION NINETY-ONE NINETY-ONE

SECTION NINETY-TWO NINETY-TWO

SECTION NINETY-THREE NINETY-THREE

SECTION NINETY-FOUR NINETY-FOUR

SECTION NINETY-FIVE NINETY-FIVE

SECTION NINETY-SIX NINETY-SIX

SECTION NINETY-SEVEN NINETY-SEVEN

SECTION NINETY-EIGHT NINETY-EIGHT

SECTION NINETY-NINE NINETY-NINE

SECTION ONE HUNDRED ONE HUNDRED

SECTION ONE HUNDRED-ONE ONE HUNDRED-ONE

SECTION ONE HUNDRED-TWO ONE HUNDRED-TWO

SECTION ONE HUNDRED-THREE ONE HUNDRED-THREE

SECTION ONE HUNDRED-FOUR ONE HUNDRED-FOUR

SECTION ONE HUNDRED-FIVE ONE HUNDRED-FIVE

SECTION ONE HUNDRED-SIX ONE HUNDRED-SIX

SECTION ONE HUNDRED-SEVEN ONE HUNDRED-SEVEN

SECTION ONE HUNDRED-EIGHT ONE HUNDRED-EIGHT

SECTION ONE HUNDRED-NINE ONE HUNDRED-NINE

SECTION TWO HUNDRED TWO HUNDRED

SECTION TWO HUNDRED-ONE TWO HUNDRED-ONE

SECTION TWO HUNDRED-TWO TWO HUNDRED-TWO

SECTION TWO HUNDRED-THREE TWO HUNDRED-THREE

SECTION TWO HUNDRED-FOUR TWO HUNDRED-FOUR

SECTION TWO HUNDRED-FIVE TWO HUNDRED-FIVE

SECTION TWO HUNDRED-SIX TWO HUNDRED-SIX

SECTION TWO HUNDRED-SEVEN TWO HUNDRED-SEVEN

SECTION TWO HUNDRED-EIGHT TWO HUNDRED-EIGHT

SECTION TWO HUNDRED-NINE TWO HUNDRED-NINE

SECTION THREE HUNDRED THREE HUNDRED

SECTION THREE HUNDRED-ONE THREE HUNDRED-ONE

SECTION THREE HUNDRED-TWO THREE HUNDRED-TWO

SECTION THREE HUNDRED-THREE THREE HUNDRED-THREE

SECTION THREE HUNDRED-FOUR THREE HUNDRED-FOUR

SECTION THREE HUNDRED-FIVE THREE HUNDRED-FIVE

SECTION THREE HUNDRED-SIX THREE HUNDRED-SIX

SECTION THREE HUNDRED-SEVEN THREE HUNDRED-SEVEN

SECTION THREE HUNDRED-EIGHT THREE HUNDRED-EIGHT

SECTION THREE HUNDRED-NINE THREE HUNDRED-NINE

SECTION FOUR HUNDRED FOUR HUNDRED

SECTION FOUR HUNDRED-ONE FOUR HUNDRED-ONE

SECTION FOUR HUNDRED-TWO FOUR HUNDRED-TWO

SECTION FOUR HUNDRED-THREE FOUR HUNDRED-THREE

SECTION FOUR HUNDRED-FOUR FOUR HUNDRED-FOUR

SECTION FOUR HUNDRED-FIVE FOUR HUNDRED-FIVE

SECTION FOUR HUNDRED-SIX FOUR HUNDRED-SIX

SECTION FOUR HUNDRED-SEVEN FOUR HUNDRED-SEVEN

SECTION FOUR HUNDRED-EIGHT FOUR HUNDRED-EIGHT

SECTION FOUR HUNDRED-NINE FOUR HUNDRED-NINE

SECTION FIVE HUNDRED FIVE HUNDRED

SECTION FIVE HUNDRED-ONE FIVE HUNDRED-ONE

SECTION FIVE HUNDRED-TWO FIVE HUNDRED-TWO

SECTION FIVE HUNDRED-THREE FIVE HUNDRED-THREE

SECTION FIVE HUNDRED-FOUR FIVE HUNDRED-FOUR

SECTION FIVE HUNDRED-FIVE FIVE HUNDRED-FIVE

SECTION FIVE HUNDRED-SIX FIVE HUNDRED-SIX

SECTION FIVE HUNDRED-SEVEN FIVE HUNDRED-SEVEN

SECTION FIVE HUNDRED-EIGHT FIVE HUNDRED-EIGHT

SECTION FIVE HUNDRED-NINE FIVE HUNDRED-NINE

SECTION SIX HUNDRED SIX HUNDRED

SECTION SIX HUNDRED-ONE SIX HUNDRED-ONE

SECTION SIX HUNDRED-TWO SIX HUNDRED-TWO

SECTION SIX HUNDRED-THREE SIX HUNDRED-THREE

SECTION SIX HUNDRED-FOUR SIX HUNDRED-FOUR

SECTION SIX HUNDRED-FIVE SIX HUNDRED-FIVE

SECTION SIX HUNDRED-SIX SIX HUNDRED-SIX

SECTION SIX HUNDRED-SEVEN SIX HUNDRED-SEVEN

SECTION SIX HUNDRED-EIGHT SIX HUNDRED-EIGHT

SECTION SIX HUNDRED-NINE SIX HUNDRED-NINE

SECTION SEVEN HUNDRED SEVEN HUNDRED

SECTION SEVEN HUNDRED-ONE SEVEN HUNDRED-ONE

SECTION SEVEN HUNDRED-TWO SEVEN HUNDRED-TWO

SECTION SEVEN HUNDRED-THREE SEVEN HUNDRED-THREE

SECTION SEVEN HUNDRED-FOUR SEVEN HUNDRED-FOUR

SECTION SEVEN HUNDRED-FIVE SEVEN HUNDRED-FIVE

SECTION SEVEN HUNDRED-SIX SEVEN HUNDRED-SIX

SECTION SEVEN HUNDRED-SEVEN SEVEN HUNDRED-SEVEN

SECTION SEVEN HUNDRED-EIGHT SEVEN HUNDRED-EIGHT

SECTION SEVEN HUNDRED-NINE SEVEN HUNDRED-NINE

SECTION EIGHT HUNDRED EIGHT HUNDRED

SECTION EIGHT HUNDRED-ONE EIGHT HUNDRED-ONE

SECTION EIGHT HUNDRED-TWO EIGHT HUNDRED-TWO

SECTION EIGHT HUNDRED-THREE EIGHT HUNDRED-THREE

SECTION EIGHT HUNDRED-FOUR EIGHT HUNDRED-FOUR

SECTION EIGHT HUNDRED-FIVE EIGHT HUNDRED-FIVE

SECTION EIGHT HUNDRED-SIX EIGHT HUNDRED-SIX

SECTION EIGHT HUNDRED-SEVEN EIGHT HUNDRED-SEVEN

SECTION EIGHT HUNDRED-EIGHT EIGHT HUNDRED-EIGHT

SECTION EIGHT HUNDRED-NINE EIGHT HUNDRED-NINE

SECTION NINE HUNDRED NINE HUNDRED

SECTION NINE HUNDRED-ONE NINE HUNDRED-ONE

SECTION NINE HUNDRED-TWO NINE HUNDRED-TWO

SECTION NINE HUNDRED-THREE NINE HUNDRED-THREE

SECTION NINE HUNDRED-FOUR NINE HUNDRED-FOUR

SECTION NINE HUNDRED-FIVE NINE HUNDRED-FIVE

SECTION NINE HUNDRED-SIX NINE HUNDRED-SIX

SECTION NINE HUNDRED-SEVEN NINE HUNDRED-SEVEN

SECTION NINE HUNDRED-EIGHT NINE HUNDRED-EIGHT

SECTION NINE HUNDRED-NINE NINE HUNDRED-NINE

SECTION ONE THOUSAND ONE THOUSAND

SECTION ONE THOUSAND-ONE ONE THOUSAND-ONE

SECTION ONE THOUSAND-TWO ONE THOUSAND-TWO

SECTION ONE THOUSAND-THREE ONE THOUSAND-THREE

SECTION ONE THOUSAND-FOUR ONE THOUSAND-FOUR

SECTION ONE THOUSAND-FIVE ONE THOUSAND-FIVE

SECTION ONE THOUSAND-SIX ONE THOUSAND-SIX

SECTION ONE THOUSAND-SEVEN ONE THOUSAND-SEVEN

SECTION ONE THOUSAND-EIGHT ONE THOUSAND-EIGHT

SECTION ONE THOUSAND-NINE ONE THOUSAND-NINE

PROCESSING AND PREPARATION NOTES																									
<p>Simple method for the measurement of dichroism. N. M. Melancholia. <i>Soviet Geol</i> 1940, Nos. 5-6, 184-7. Using a synthetic ruby as an example, M compares the dichroism as measured by the spectrophotometric and the dichroscopic ocular methods. The former gives a slightly higher max. at $\lambda = 560$ mμ, shifted very slightly to longer wave lengths. F. H. Rathmann</p>																									
<p>ASB SLA METALLURGICAL LITERATURE CLASSIFICATION</p>																									

CA

8

The absorption curve of iron oxide in minerals. N. M. Melnikholin (Inst. Mineral Raw Materials, Moscow) *Zapiski Vserossiiskogo Mineral. Obshchestva* (Mém. soc. russe minéral.) 75, 89-93 (1946); *Chem. Zentr.* (Russian Zone Ed.) 1948, I, 527. — Measurements were made of the absorption of hematite inclusions in muscovite, red sylvite, the pigment formed from kaolin and Fe_2O_3 , artificial carnelian, and sphalerite. The results showed good agreement, the curves showing 3 narrow, low max. at about 470, 490, and 515 m μ . In yellow corundum these were displaced somewhat downward. Epidote and green demantoid showed a sharp max. at about 430 m μ .

M. G. Moore

CA

1

New model of a polarization interferometer for the microscope. N. M. Melanikhin. *Zavodskaya Lab.* 13, 495-7(1947).--An app. involving 3 quartz plates cut at 45° to the optical axis, and a half-wave plate, can be attached to any microscope for the detn. of refractive indexes. N. Thon

MELANKHOLIN, N.M., inzhener.

Optical method of controlling the thickness of mica sheets. Vest.elektro-
prom. 18 no.6:23-24 Je '47. (MLRA 6:12)

1. Institut kristallografii Akademii nauk SSSR.

(Mica)

MELANKHOLIN, N.M.

The coloration of nicas. Trudy Inst. Krist., Akad. Nauk S.S.S.R.
4, 223-9 '48.
(CA 47 no.13:6313 '53)

MELANKHOLIN, N. M.

Refraction index measurement under microscope by immersion method. Moskva, Izd-vo Akad. nauk S.S.R., 1949. 99 p. (51-21030)

QE4 .3.5.M4

MELANKHOLIN, N. M.

Refractive Index

Application of new model of polarization interferometer for measuring indices of refraction of immersion media. Trudy Inst. krist., No. 5, 1949.

Monthly List of Russian Accessions, Library of Congress, December 1952. Unclassified.

MELANKHOLIN, N. M.

PA 3/50739

USSR/Geology - Coal
Carbon, in Coal

Sep/Oct 49

"Determining the Degree of Carbonification (Coal Rank) of Mineral Coals by Measuring Their Elliptical Polarization Constant," N. M. Melankholin, 8 pp

"Iz Ak Nauk SSSR, Ser Geol" No 5

Degree of carbonification, quantitatively characterized by content of volatile components and carbon in coal, determines commercial grade of coal. Attempted to establish relation between degree of carbonification and optical constants of coals, particularly constants of elliptical polarization (angles of polarization), for specimens of coals of various grades from the Donetsk Basin. To measure these constants, a special instrument was made by adding a number of devices to a single-circle gonimeter. Results of measurements showed a definite relationship between values of the constant and content of carbon and volatile components in the coal. Established dependence of the constant upon petrographic composition of coals. Due to unavoidable emergence of thin-adsorption films on surface of coals, measured values for elliptical polarization constant are not absolutely correct. However, they can be used successfully to determine grade

3/50739

USSR/Geology - Coal (Contd 2)

Sep/Oct 49

of coal, thus replacing chemical analysis by optical analysis.

3/50739

1. MELANKHOLIN, N. M.; RUDNITSKAYA, YE. S.
2. USSR 600
4. Quartz
7. Optical properties of a Rochon quartz prism, Trudy Inst. krist, No. 7, 1952.
9. Monthly List of Russian Accessions, Library of Congress, April 1953, Uncl.

1. MELANKHOLYN, N. M.
2. USSR (600)
4. Crystallography
7. Some properties of an obliquely orientated quartz wedge. Trudy Inst.krist., no. 7, 1952.
9. Monthly List of Russian Accessions, Library of Congress, April 1953, Uncl.

MELANEKHOLIN, N. M.

Crystallography

A description of a series of compensations to obliquely
oriented cuts of minerals. Zap. Vses. min. ob.
81 no. 1:54-55 '52

Monthly List of Russian Accessions, Library of
Congress, July 1952. Unclassified

MELANKHOLIN, N.M.

Measurement of the refractive index of synthetic corundum and the
corundum batch. Trudy Inst.krist. no.8:57-76 '53. (MLRA 7:5)
(Corundum) (Refractive index)

MELANKHOLIN, N.M.

Use of Tsepler's method for the study of nonhomogeneity in synthetic
corundum crystals. Trudy Inst.krist. no.8:189-192 '53. (MLRA 7:5)
(Corundum)

MELANKHOLIN, N. M.

PHASE I TREASURE ISLAND BIBLIOGRAPHICAL REPORT

AID 468 - I

BOOK

Call No.: AF637791

Authors: MELANKHOLIN, N. M. and GRUM-GRZHIMAYLO, S. V.

Full Title: METHODS OF INVESTIGATING THE OPTICAL PROPERTIES OF CRYSTALS

Transliterated Title: Metody issledovaniya opticheskikh svoystv kristallov

PUBLISHING DATA

Originating Agency: Academy of Sciences, USSR. Institute of Crystallography

Publishing House: Academy of Sciences, USSR

Date: 1954 No. pp.: 192

No. of copies: 3,500

Editorial Staff

Editor: Vedeneyeva, N. Ye., Doctor of Phys. and Math. Sci.

Others: The staff of the Institute of Crystallography of the Ac. of Sci., USSR, and the staff of the crystallography laboratory previously with the All-Union Scientific Institute for Research in Minerals.

PURPOSE: This is a textbook dealing with optical methods for investigating crystals and for measuring their optical constants. It is intended mainly for workers in scientific research institutes and factory laboratories investigating crystals and all kinds of crystalline products, raw material and semi-products. It can also be used as a textbook for petrographic studies.

1/2

Metody issledovaniya opticheskikh svoystv kristallov

AID 468 - I

TEXT DATA

Coverage: This is the latest Soviet textbook on optical crystallography.

It covers: methods of obtaining polarized light (polarizers); the polarizing microscope; measurements of the refractive indexes of crystals (immersion and other methods); measurement of the rotating plane of polarization of light in crystals; investigation of the absorption of light by crystals (color and pleochroism of crystals); and orientation of crystals and their investigation in converging and parallel polarized light by means of special apparatuses. The methods and apparatuses outlined are those applied to transparent, colorless and slightly absorbing crystals, which can be investigated with transparent light. Thus, methods of investigation of crystals with metallic absorption which require the study of reflected light are not covered in this book.

No. of References: 22, Russian 17 (1913-1952), Foreign 5 (1914-1950)

Facilities: None

2/2

USSR/Physics - MELANKHOLIN, N. M.
Relaxation, Measuring Instruments

FD 367

Card 1/1

Author : Regal', V. R. and Melankholin, N. M.

Title : Rigid optical dynamometer

Periodical : Zhur. tekhn. fiz. 24, 454-459, Mar 1954

Abstract : Describes dynamometer for measuring stress relaxation in compressed specimens. Compressing force is measured by determination of double refraction in a crystal-measurer which is made of a single crystal of TlI-TlBr solid solution. Design is based on sufficiently high rigidity of crystal-measurer, so its deformation is considerably lower than that of test specimen. Instrument, being equipped with special optical system, is used in combination with microscope and serves for plotting relaxation characteristics of various materials. One USSR reference since 1953. Illustrations.

Institution :

Submitted : October 16, 1953

Category : USSR/Optics - Physical Optics

K-5

Abs Jour : Ref Zhur - Fizika. No. 1957, No. 2310

Author : Melankholia, N M.

Title : New Instrument for Optical Orientation of a Quartz Pebble

Orig Pub : Tr. In-ta kristallogr. AN SSSR 1965 vyp. 11. 239-242

Abstract : Description of a simple instrument for the determination of the orientation of the optical axis of a quartz pebble; this axis coincides with the principal crystallographic axis of the quartz crystal. The direction of the principal axis is first determined approximately, and then a slab, approximately parallel to the optical axis, is cut from the pebble. The arrangement of the instrument permits rotation of such a slab in its plane, inclination of the slab about its parallel axis, and simultaneous observation of the conoscopic figure that is directly visible to the eye when the slab is located between crossed polaroids. By mounting the optical axis of the slab vertically, it is thus possible to measure the direction and the angle of inclination of the axis relative to the normal to the slab with an accuracy to 0.25° . From these data it is possible to obtain a cut on the tested pebble perpendicular to the optical axis, and to determine the directions of the electric axis on the pebble. The same instrument can also be used to determine the sign of the rotation of the plane of polarization in the tested pebble, by rotating the upper polaroid or by introducing an additional quartz plate.

Card : 1/1

USSR/Physics - Crystals disruption

FD-3252

Card 1/1 Pub. 146 - 11/44

Author : Melankholin, N. M.; Regel', V. R.

Title : Experience gained in an investigation of the process of disruption of NaCl crystals

Periodical : Zhur. eksp. i teor. fiz., 29, No 6(12), Dec 1955, 817-821

Abstract : An investigation, by the method of cinematography under polarized light, of the process governing the development of fissures during stretching of specimens of NaCl single-crystals. It is shown that the fissures are propagated in jump-like fashion. The development of fissures is prevented by the shear bands, which must be intersected by the fissures. The authors assume that the prevention of the propagation of fissures is caused by those regions of the shear bands in which compressing stresses exist. The regions of the shear bands with tensile stresses evidently must not prevent the development of fissures. The authors thank Professor M. V. Klassen-Neklyudova for her advice. Twelve references: e.g. M. V. Klassen-Neklyudova, *ibid.*, 6, 584, 1936.

Institution : Institute of Crystallography, Academy of Sciences USSR

Submitted : June 14, 1954


MELENKHOLOV, N. M.

Experiments for the study of gilding phenomena in sodium chloride crystals. N. M. Melankholla and V. B. Regel. *Trudy Inst. Krist., Akad. Nauk SSSR*, 1936, No. 12, 148-51. The previous micro-cinematographic results of Stepanov and Mil'kamanovich (cf. *C.A.* 45, 9975ag, 9976c) are confirmed by improved exptl. methods in polarized light between crossed Nicols. The problem of holding the crystal at high temps. was considered carefully. The relative local distribution of compressive and tensile strains in the crystal that induce the shear gilding translation phenomena are described in detail. Under a constantlv increasing load the...

2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100

"APPROVED FOR RELEASE: 07/12/2001

CIA-RDP86-00513R001033330003-5



APPROVED FOR RELEASE: 07/12/2001

CIA-RDP86-00513R001033330003-5"

Microcrystals 40

USSR/Optics -Optical Methods of Analysis. Instruments.

K-7

Abs Jour : Referat Zhur - Fizika, No 5, 1957, 13135

Author : Melankholin, N.M., Grechushnikov, B.N.

Inst : Institute of Crystallography, Academy of Sciences, USSR

Title : New Photoelectric Microspectrophotometer.

Orig Pub : Tr. in-ta Pristallogr. AN SSSR, 1956, vyp. 12, 186-191

Abstract : To investigate the absorption spectra of various small crystals of organic dyes (up to 0.03 mm in diameter), a photoelectric microspectrophotometer was constructed. The source of monochromatic light is the UM-2 monochromator, and the receiver is a FEU-19 photomultiplier. The darkness current of the photomultiplier is compensated for. The instrument can be used for the investigation of the absorption spectra of all microcrystals or biological objects, and also for the spectrophotometric

Card 1/2

Abs Jour : Ref Zhur - Fizika, No 5, 1957, 13135

MELANKHOLIN, N. M.

Handwritten: $D = \log I_0/I$

Handwritten: 1/2

Colors of amphiboles. N. M. Melankholin. *Zapiski Vsesoyuz. Mineralog. Obshchestva* 85, 218-22 (1956). The absorption curves for a series of amphiboles were detd. with a photoelec. spectrophotometer, and the photometer eyepiece was adapted to a polarization microscope (cf. Melankholin (1940)). The ordinates of the curves are the optical ds. $D = \log I_0/I$, as a function of the wave length of the visible light. The measurements were made for the principal indicatrix axes, γ and α , also the pleochroism was detd. The investigated amphiboles were: green and brown hornblende, basaltic hornblende, edenite, arfvedsonite, glaucophane, and actinolite, chiefly from mineral deposits of the Ukraine and Kamchatka. The curves show in most cases distinctly the superposition of the characteristic absorption of the Fe^{2+} and Fe^{3+} cations (cf. Koltic, *C.A.* 29, 5777e; Melankholin, *C.A.* 44, 7717e). Specific effects of the compn. are discussed for the extinction curves of glaucophane in which the Na^+ cations exert some polarization on the O^{2-} anions surrounding the Fe^{2+} color centers. A deep min. of absorption is characteristic at 500 $m\mu$ and a distinct shifting of the absorption max. in infrared and red to the violet end of the visible spectrum. The curves for arfvedsonite are similar, although the max. in red is not so strongly shifted as in glaucophane; the anisotropy of the absorption is more distinct than in glaucophane. The transition of the absorption phenomena in green to brown hornblendes is continuous; the curves for the green hornblendes are rather similar to those of arfvedsonite. The higher the contents in Fe^{2+} the more the typical curves of brown hornblende are observed, with a rather flat min. above 600 $m\mu$. In the deep-brown basaltic hornblendes this min. is in the infrared, and the curves for γ and α absorption are sybatic. Actinolite shows only very flat curves for α and γ , without distinct max. or min. Al^{3+} does not affect the character of the absorption curves for the Fe-contg. amphiboles. The relative

MELANKHOLIN, N. M.

"distances between the α and γ curves are a measure for the degree of dichroism; they are different in glaucophane-actinolite, and the hornblende-amphiboles. In the first group the dichroism is stronger in the red; in basaltic it is stronger in the green-blue portions of the spectrum, in green hornblendes it is intermediate. Evidently, the position of the Fe^{2+} cations in groups $[\text{FeO}_4]$ is different from that of the Fe^{3+} cations in the structure of the amphiboles, with respect to the $[\text{Si}_2\text{O}_6]$ ribbons.

W. Rittel

2/2

Melankholin, N. M.
USSR/Physical Chemistry. Molecules. Chemical bond.

B-4

Abs Jour : Ref Zhur - Khimiya No 7, 1957, 21979

Author : Melankholin, N. M.

Inst : None

Title : Absorption spectra of methylene-blue crystals.

Orig Pub : Dokl. AN SSSR 1956, 108, No 2, 247-250

Abstract : The absorption spectra of microcrystals of methylene blue (up to 20 μ in diameter) were investigated in polarized light. Because of the different orientation of monoclinic crystals of this dyestuff, which are formed on glass when a drop of solution is evaporated, the spectrum components with polarization of incident light along all three axis of indices were obtained. The highest absorption was found when the light was polarized along the axis of largest refraction index, which coincides with the crystalline axis. The absorption band has a maximum around 550m μ and corresponds to band around 656m μ in the spectrum of solution. When the light was polarized along the axis of smallest light refraction a maximum around 620m μ was found. The absorption curve for light vibration parallel to the middle refraction index has a maximum

Card 1/2

- 9-

USSR/Physical Chemistry. Molecule. Chemical bond.

B-4

Abs Jour : Ref Zhur - Khimiya No 7, 1957, 21979

around $405\text{m}\mu$. Besides the absorption bands for vibrations directed along the crystalline axis, there is another band with maximum around $490\text{m}\mu$, corresponding to a direction of vibration, which is under an angle to the axis. Presence of this band, according to the author, indicates that the form of area of absorption of crystal is not shaped like an ellipse or oval, but resembles the area of crystal's elasticity. The author associates the $550\text{m}\mu$ band with molecular absorption of the dyestuff, calling the other 3 absorption bands - 620, 490 and $405\text{m}\mu$ - pure crystalline ones.

Card 2/2

-10-

PRIKHOTKO, A F
24(7) 3 PHASE I BOOK EXPLOITATION SOV/1365
L'vov. Universitet
Materialy I Vsesoyuznogo soveshchaniya po spektroskopii. t. 1:
Molekulyarnaya spektroskopiya (Papers of the 10th All-Union
Conference on Spectroscopy. Vol. 1: Molecular Spectroscopy)
[L'vov] Izd-vo L'vovskogo univ-ta, 1957. 499 p. 4,000 copies
printed. (Series: Its: Fizichnyy zbirnyk, vyp. 3/8/)
Additional Sponsoring Agency: Akademiya nauk SSSR. Komissiya po
spektroskopii. Ed.: Jazer, S.L.; Tech. Ed.: Saranyuk, T.V.;
Editorial Board: Lavsterg, G.S., Academician (Resp. Ed., Deceased),
Neporent, B.S., Doctor of Physical and Mathematical Sciences,
Fabelinskiy, I.L., Doctor of Physical and Mathematical Sciences,
Fabrikant, V.A., Doctor of Physical and Mathematical Sciences,
Kornitskiy, V.G., Candidate of Technical Sciences, Rayskiy, S.M.,
Candidate of Physical and Mathematical Sciences, Klimovskiy, L.K.,
Candidate of Physical and Mathematical Sciences, Miliyanchuk, V.S.,
A. Ye., Candidate of Physical and Mathematical Sciences.
Card 1/30

Shatalov, A.A. Spectral Study of the Colloidal
Coagulation of F-centers in Alkali Halide Crystals 148
Pialkovskaya, O.V. Infrared Absorption Spectra of
Anthracene 151
Vartanyan, A.T. Absorption Spectra of Sublimated Dye
Layers 154
Melankholin, N.M. Absorption Spectra of Thiazine-dye
Crystals 157
Fribytkova, N.N., and L.S. Agroskin. Study of the
Optical Properties of Some Dyes in Large Samples
by the Method of Mirror Reflection 158
Zhidkova, Z.V., and Yu. M. Suss. Study of the Effect
of the Degree of Dispersion and Nature of the Ad-
sorbent on the Spectral Absorption Curve of Absorbed
Sensitized Dyes 161
Card 11/30

MELANKHOLIN, N. M.

Anomalous dispersion of the bisectrices of organic dye crystals. N. M. Melankholin. Kristallografiya 2, 78-82 (1957). Measurements of the dispersion of the bisectrices in crystals of methylene green, methylene blue, and thionine blue have revealed its anomalous character.

...the dispersion of the direc-
tions in crystals of methylene green, methylene blue,
and thionine blue have revealed its anomalous character
connected with the absorption bands of the dyes. Max.
dispersion was found in crystals with oblique orientation.
E. Ryshkevitch //

MELANKHOLIN, N. M.

51-2-2/15

AUTHOR: Melankholin, N.M.


TITLE: Absorption spectra of thiazine-dye crystals. (Spektry pogloshcheniya kristallov tiazinovykh krasiteley).

PERIODICAL: "Optika i Spektroskopiya" (Optics and Spectroscopy) 1957, Vol.3, No.2, pp.104-114 (U.S.S.R.)

ABSTRACT: Study of absorption spectra in the range 400-650 m μ of three organic dyes: methylene blue, new methylene blue and methylene green. This work is an extension of the investigation of absorption of monocrystals of the above three dyes by N.E.Vedeneeva and E.N.Slavnova (2) (the data obtained by these two authors were not very precise). A microspectrophotometer consisting of a monochromator, a polarizing microscope and a photomultiplier was used. Since strongly pleochroic crystals should be studied in nearly parallel light, diaphragms were used at the microscope objective and at the condenser so that angle of convergence of light was about 5-6° in the crystal (placed on the object stage of the microscope). The crystals were prepared by slow evaporation of an aqueous solution of the dye placed on an object slide. On one slide crystals of various habits and orientations were obtained.

Card 1/6

Absorption spectra of thiazine-dye crystals. (Cont.)^{51-2-2/15}



crystallizes as long prisms with their c-axis approximately parallel to the slide surface. In addition to crystals oriented films of methylene blue (Demon (4)) were studied. Absorption curves for vibrations parallel to the N_m (i.e. almost parallel to the c-axis) and N_p (i.e. almost parallel to the a-axis) axes of the optical indicatrix were found using monocrystals. Since the vibrations parallel to the N_g indicatrix axis (and the crystal b-axis are strongly absorbed, oriented films were used. These films, prepared by rolling a glass rod wetted with dye solution across a heated slide, consist of similarly oriented crystallites whose a and N_p axes are perpendicular to the glass slide. The absorption curves obtained are in Figs. 1, 2 and 3. For the vibrations parallel to N_g a strong absorption maximum occurs at about $550 \text{ m}\mu$ (Fig.1). A gentle maximum is observed around $615 \text{ m}\mu$ for the vibrations parallel to the N_p axis (Fig.2). A maximum at $405 \text{ m}\mu$ is found for the vibrations parallel to N_m (Fig.2). A further


Card 2/6

51-2-2/15

Absorption spectra of thiazine-dye crystals. (Cont.)

maximum (Fig.3) at $490 \text{ m}\mu$ corresponds to vibrations which are in the N_g-N_m plane but lie in two directions symmetrical with respect to the N_g and the N_m axes. The most intense absorption band at $550 \text{ m}\mu$ is a strongly displaced molecular band and the others are due to intermolecular bonds in the crystal.

New methylene blue.



results are given in Figs.4, 5 and 6. It was not possible to assign the absorption curves to crystallographic directions, since the crystal structure of new methylene blue (and methylene green) is not yet known. The absorption maxima occur at about $640 \text{ m}\mu$ for the vibrations parallel to N_g (Fig.4) and at $570 \text{ m}\mu$ for N_m (Fig.5). For the vibrations parallel to N_p two repeatable sets of maxima are obtained: at $595 \text{ m}\mu$ for the N_p-N_m crystal cut (Fig.5) and at $605 \text{ m}\mu$ for the N_g-N_p cut (Fig.4). Fig.6 shows the results for directions which are

Card 3/6

51-2-2/15

Absorption spectra of thiazine-dye crystals. (Cont.)

assigned to vibrations oblique with respect to the indicatrix axes. Fig.9 shows absorption curves for a cut near to the N_g-N_m plane. The results obtained indicate that absorption by the dye crystals studied here can be due to molecular and crystalline (i.e. intermolecular bonds in crystals) properties. When the crystal structure is known (methylene blue) the molecular bands can be identified. A characteristic property of the crystals studied is the occurrence of absorption bands corresponding to vibrations oriented obliquely with respect to the indicatrix axes. All the three dyes exhibit strong absorption bands at $550-570\text{ m}\mu$ for one of the vibration directions perpendicular to the longitudinal axis of the crystal. For both (1) methylene blue and (2) new methylene blue, the absorption bands are strongly polarized; the absorption band for vibrations parallel to the short axis of the molecule lies in the short-wave region for the former and in the long-wave region for the latter. The methylene green (3) bands are weakly polarized. The differences between (1) and (2) on one hand and (3) on the other are due to different molecular symmetries. (1) and (2) possess a second-order symmetry axis along the

Card 5/6

Absorption spectra of thiazine-dye crystals. (Cont.)

51-2-2/15

short axis of the molecule through N and S atoms, while in (3) the symmetry is affected by the presence of NO_2 . There are ten figures and ten references (eight of which are Slavic). References cited: (2) and (4).

SUBMITTED: December 3, 1956.

ASSOCIATION: Crystallographical Institute, Academy of Sciences of the U.S.S.R.

AVAILABLE: Library of Congress

Card 6/6

SOV/51-5-4-12/21

AUTHOR: Melankholin, N.M.

TITLE: Absorption Spectra of Solid Films of Organic Dyes (Spektry pogloshcheniya tverdykh plenok organicheskikh krasiteley)

PERIODICAL: Optika i Spektroskopiya, 1958, Vol 5, Nr 4, pp 435-439 (USSR)

ABSTRACT: Because of the small dimensions and non-uniformities of films the author constructed a special very sensitive photoelectric microspectrophotometer with a photoelectric multiplier. This microspectrophotometer was used to obtain the absorption spectra of film areas 0.02 mm in diameter. All the films studied were prepared by slow evaporation of an aqueous solution on a microscope object slide. Oriented films of methylene blue were prepared by Demon's method (Ref 6) by rolling a glass rod wetted with a dye solution across a heated glass slide. Certain of the dyes studied, such as thionine violet produce only crystalline films, other dyes (e.g. thionine blue) produce only amorphous layers which become crystalline after a certain time. Methylene blue layers may be either crystalline or amorphous. The absorption curves of methylene blue are shown in Figs 1 and 2, those of methylene green in Fig 3, and of thionine dyes in Fig 4. In Figs 1, 2 and 4 the letters "a" and "b" refer to measurements at two mutually perpendicular directions of polarized light. Oriented

Card 1/2

Absorption Spectra of Solid Films of Organic Dyes

SOV/51-5-4-12/21

methylene blue layers were found to be crystalline (Fig 1, curves 1, 2; Fig 2, curves 2), but amorphous layers, blue in colour, could be also obtained (Fig 2, curves 1). Methylene green layers (Fig 3, curve 1 represents blue layers and Fig 3, curve 2 represents rose layers) were obtained in the oriented crystalline form. Thionine violet (Fig 4, curves 1) and toluidine blue form crystalline layers, which are not necessarily oriented. Thionine blue (Fig 4, curve 2) and Capri blue form amorphous layers. The crystalline layers consist of microcrystallites of various forms, which may be oriented along one direction. The amorphous layers consist of disordered aggregates of dimer molecules of a dye and of water molecules. There are 4 figures, and 7 references, 6 of which are Soviet and one American.

ASSOCIATION: Institut Kristallografii, AN SSSR (Institute of Crystallography, Academy of Sciences of the U.S.S.R.)

SUBMITTED: October 28, 1957.

Card 2/2

1. Dye films--Spectra 2. Microspectrophotometers--Applications

Melankhola, V. M.

24(2)

PHASE I BOOK EXPLOITATION

SOV/2353

Akademiya nauk SSSR. Institut kristallografi

Rost kristallov, tom. 2 (Growth of Crystals, Vol. 2) Moscow, 1959. 238 p.
Errata slip inserted. 2,000 copies printed.

Resp. Eds.: A. V. Shubnikov, Academician, and N. N. Sheftal', Doctor of
Geological and Mineralogical Sciences; Ed. of Publishing House:
K. S. Aleksandrov; Tech. Ed.: T. V. Polyakova.

PURPOSE: This book is intended for scientists and researchers engaged in
crystallography and in growing industrial monocrystals.

COVERAGE: This is the second of two volumes on crystal growth. The first
volume contained reports delivered at the First Congress on Crystal Growth.
The present volume also contains an extensive study of corundum synthesis
by S. K. Popov [deceased]. These studies reflect the development of Soviet
research in crystallography in the period following the first congress.
The studies contain some essentially new results obtained by Soviet scientists.
The editors express the hope that these studies will unite the efforts of Sov-
iet scientists engaged in studying the process of crystal growth and in grow-

Card 1/5

Growth of Crystals (Cont.)

SOV/2353

ing industrially valuable monocrystals. No personalities are mentioned.
References are given at the end of each article.

TABLE OF CONTENTS:

Foreword

3

I. THEORETICAL AND EXPERIMENTAL STUDIES

Khaimov-Mal'kov, V. Ya. Thermodynamics of Crystallization Pressure

5

Khaimov-Mal'kov, V. Ya. Problem of Experimental Determination of the
Crystallization Pressure Value

17

Khaimov-Mal'kov, V. Ya. Growth Conditions of Crystals in Contact
With a Macroobstruction

26

Kapustin, A. P., and V. Ye. Kovalyunayte. Crystallization of Alum
From Aqueous Solutions in an Ultrasonic Field

40

Card 2/5

Growth of Crystals (Cont.)

SOV/2353

Slavnova, Ye. N. Adsorption of Thionine Blue by Growing Crystals of Lead Nitrate and Its Effect on the Morphological Properties of the Crystals 44

Melankholin, N. M. Growth Forms of Crystals of Thiazine Dyes 49

Tsinober, L.I., L. G. Chentsova, and A. A. Shternberg. The Green and Brown Tint of Synthetic Quartz Crystals 61

Kamentsev, I. Ye. Crystallization of KJ on a Biotite and Muscovite Surface 68

Belyustin, A. V. Possibility of Determining Surface Energy of Crystals From Equilibrium States 70

II. GROWING OF MONOCRYSTALS (APPARATUS, METHODS, SUPPLEMENTARY WORK)

Kashpar, Yan. Growing of Calcite and of Other Carbonates 73

Spendiarov, N. N., and B. S. Aleksandrov. Growing of Anthracene Monocrystals 78

Card 3/ 5

SOV/70-4-4-15/34

AUTHORS: Melankholin, N.M. and Slavnova, Ye.N.

TITLE: Regularities in the Take-up of Capri Blue By Growing Crystals of Lead Nitrate

PERIODICAL: Kristallografiya, 1959, Vol 4, Nr 4, pp 563-570 (USSR)

ABSTRACT: A spectrophotometric method was used to study the capture of Capri blue (as illustrated on p 563) by growing crystals of $Pb(NO_3)_2$ and to construct the corresponding adsorption curve. The appearance of the crystals depended on the concentration of dye in the solution. It was shown that the dye exists in the lead nitrate crystals as sub-microscopic crystallites. Crystal-optical studies of the lead nitrate crystals disclosed different physical properties for the positive and negative tetrahedra. Lead nitrate crystals were grown at room temperature with dye concentrations of 2.0×10^{-3} to 0.50 weight % : Dye is only adsorbed when its concentration is $> 10^{-2}$ %. Concentrations of up to 10^{-1} % crystals grow as cubotetrahedra and for higher concentrations as cubes. The colouring of all faces is not the same and the growth pyramids of the cube faces are

Card1/2

SOV/70-4-4-15/34

Regularities in the Take-up of Capri Blue By Growing Crystals of Lead Nitrate

intensely coloured even at small concentrations. The absorption spectrum of the dye as adsorbed corresponds neither to that of the monomer nor to that of the dimer molecules but to that of the crystalline material. The orientation of the molecules can be deduced from the dichroism. This orientation is the same in all parts of the crystal. There are 7 figures, 1 table and 4 Soviet references.

ASSOCIATION: Institut kristallografii AN SSSR (Institute of Crystallography of the Ac.Sc., USSR)

SUBMITTED: March 17, 1959

Card2/2

24.7100

AUTHORS: Melankholin, N. M., Abramovich, S. G.

TITLE: Optical Identification of the Crystalline Modifications of Indanthrene Blue RS. Brief Communication

PERIODICAL: Kristallografiya, 1959, Vol 4, No 6, pp 900-901 (USSR)

ABSTRACT: As shown by Ye. N. Belova (Crystallographical Institute of the Academy of Sciences, USSR), the powder diffraction patterns of the known four crystalline modifications of indanthrene blue RS differ sufficiently to identify them, except in the case of γ and δ modifications whose diffraction patterns are very close to each other. By crystal optical and spectrophotometric study of the crystals and of a number of trade specimens, the authors found that dry specimens can be identified optically. As can be seen under a polarization microscope, α and δ form well-shaped crystals, while β and γ occur in amorphous-looking formless aggregates in which

Card 1/2

Optical Identification of the Crystalline
Modifications of Indanthrene Blue RC.
Brief Communication

77128
SOV/10-4-6-19/31

only a few poorly-shaped crystalline grains can be seen. Fine prismatic α has strong double refraction and weak dichroism, while the rhombic fine platelets of δ show strong pleochroism from light blue to dark violet, and weak double refraction. Absorption curves of α and γ , obtained by automatic spectrophotometer SF-2m, show definite maxima within the range of visible spectrum, i.e., α at 625 m μ and γ at 590 m μ . No clear maximum appears in the absorption curves of β and δ (Fig. 1). Combining the data of the absorption curves with those obtained by polarization microscopic study, the 4 modifications could be identified in trade specimens. There is 1 figure; and 2 references, 1 Soviet, 1 German.

Card 2/4 *Inst. Crystallography, AS USSR*
Sci. Res. Inst. Organic Intermediate Products & Dyes

SOV/51-7-4-10/32

AUTHOR: Melankholin, N.M.

TITLE: Absorption Spectra of Crystals of Certain Thiazine and Oxazine Dyes.

PERIODICAL: Optika i spektroskopiya, 1959, Vol 7, Nr 4, pp 498-504 (USSR)

ABSTRACT: The present paper is a continuation of earlier work (Refs 1-3). It reports results obtained for two thiazine dyes (thionine blue and toluidine blue) and one oxazine dye (Capri blue). Crystals of these dyes were produced by slow evaporation of drops of their aqueous solutions. The absorption curves of all these dyes were obtained by means of a specially constructed photoelectric microspectrophotometer (Ref 4). This instrument made it possible to carry out measurements on crystals of 0.02 mm dimensions in a polarized beam of 90° aperture in air. Since the thickness of the crystals used could not be measured exactly and their surfaces were poor, the absorption coefficients were not calculated and all the absorption curves are shown in the form of the dependence of the optical density on wavelength (400-800 mμ). These absorption curves are given in Figs 1-5; they represent absorption of light vibrations oriented in various ways with respect to the crystal axes. In crystals of thionine blue absorption bands at 535 and 560 mμ were observed only at oblique orientations of the light beam with respect to the principal axes of the optical indicatrix and were absent in the

Card 1/2

OV/51-7-4-10/32

Absorption Spectra of Crystals of Certain Thiazine and Oxazine Dyes

absorption spectra corresponding to the two symmetric orientations. Another peculiarity of the absorption spectrum of thionine blue was the variation of the positions of the absorption bands, corresponding to vibrations more or less parallel to the principal axes of the optical indicatrix, with the direction of the wave normal. The latter property is discussed in some detail. Apart from the absorption spectra the author studied also dispersion of the indicatrix axes in the dye crystals. Such dispersion was reported earlier for thionine blue (Ref 3). Crystals of Capri blue also had high dispersion. Measured dispersion of the extinction angle in Capri blue (Fig 6) shows that this dispersion, and consequently the dispersion of the indicatrix axes, is anomalous. There are 6 figures, 3 structural formulae and 5 Soviet references.

SUBMITTED: February 26, 1959

Card 2/2

MELANKHOLIN, N.M.

Liquid crystals of thiazine dyes. Kristallografiia 6 no.4:568-
575 JI-Ag '61. (MIRA 14:8)

1. Institut kristallografi AN SSSR.
(Liquid crystals) (Thiazine)

MELANKHOLIN, N.M.

New data on absorption spectra of thiazine dyes. Opt. 1 spektr.
11 no.2:286-289 Ag '61. (MIRA 14:8)
(Thiazine)
(Absorption spectra)

2

S/070/63/008/001/019/024
E132/E460

AUTHORS: Melankholin, N.M., Tsinober, L.I.

TITLE: The nature of biaxial crystals of synthetic amethyst

PERIODICAL: Kristallografiya, v.8, no.1, 1963, 110-112

TEXT: It has long been known that amethyst is optically biaxial, whereas quartz is uniaxial. Synthetic amethyst was made from crystals of synthetic quartz by X-ray irradiation and this was biaxial and contained Brazil twins. Plates of this material cut parallel and perpendicular to the Z-axis were examined. Most of the plates were cut from a right handed individual and in them a large number of very small left handed regions could be seen, in the form of triangular platelets, near the twin boundaries. The biaxial properties are only observed near to the twin boundaries. The angle between the optic axes varies from 0 to 7°; the optic orientation also varies and is not fixed to that of the twin boundary. It was shown that the left handed individuals in a right handed matrix do not have exactly the same orientation as the matrix differing usually by 0.7 to 0.8° and sometimes by as much as 1.2°. An explanation of the biaxial optics is given on the above basis. It can be supposed that the Brazil twins form

Card 1/2

S/070/63/008/001/019/024
E132/E460

The nature of biaxial ...

a system like a Fresnel prism and separate the right and left handed circularly polarized rays which pass along the optic axis. This is possible only on the plane perpendicular to the twin boundaries and so this explanation must be rejected. Internal strains are also rejected. A third explanation is on the basis of the difference in orientation between the left and right handed components but 1° disorientation is insufficient to account for the angle of 7° between the optic axes. The matter is not satisfactorily concluded. There is 1 figure.

ASSOCIATION: Institut kristallografii AN SSSR
(Institute of Crystallography AS USSR)

SUBMITTED: June 25, 1962

Card 2/2

ACCESSION NR: AP4009461

8/0051/63/015/006/0781/0784

AUTHOR: Melankholin, N.M.

TITLE: Absorption spectra of liquid crystals of thiazine dyes

SOURCE: Optika i spektroskopiya, v.15, no.6, 1963, 781-784

TOPIC TAGS: absorption spectrum, liquid crystal, thiazine dye, methylene blue, methylene green, novomethylene blue, thionine violet, Lauth violet, dye crystallization

ABSTRACT: In the course of an earlier investigation (N.M.Melankholin, Kristallografiya, 6, 568, 1961) in preparing samples for investigation of the absorption spectra of crystals of thiazine dyes it was discovered that there form in their water solutions liquid crystals. Microscopic investigation of the processes of crystallization showed that such liquid crystals could form in solutions of all the thiazine dyes that were available to the author. In connection with the results obtained in investigating the spectra of these dyes in true crystalline form and in water solutions, it was deemed of interest to obtain their absorption spectra in the form of liquid crystals. Accordingly, in the present study there were obtained the absorp-

Card 1/2

AP4009461

tion spectra of four thiazine dyes, namely, methylene blue, methylene green, novomethylene blue and thionine (Lauth's) violet, which form more or less stable liquid crystals of the nematic (thread-like) type. The measurements were carried out using liquid crystal films obtained by slow evaporation of drops of water solutions on glass. Uniform sections of the individual films were selected for the measurements. In view of the difficulty in determining the thickness of the films, the values of the absorption coefficient were not calculated and the curves were plotted simply on the basis of the measured values of optical density. The measurements were carried out in polarized light with the aid of the microspectrophotometer constructed by the author (Tr.Inst.kristallogr., 12, 186, 1956) for the purpose of investigating the absorption spectra of dye crystals. The absorption curves obtained for the liquid crystals are reproduced in a figure. The spectra of the individual dyes ~~are discussed in some detail and compared with the corresponding spectra of the dyes in solution and in the true crystalline state.~~ Contrary to expectation, the spectra of the liquid crystals are more similar to the spectra of the true crystals than to the spectra of solutions of the dyes. It is hypothesized that the absorption spectra of the liquid crystals are due mainly to the absorption of the crystallites composing the liquid crystals. Orig.art.has: 4 figures.

Card

2/82

Sub 25 Feb 63

MELANKHOLIN, N.M.; GUSEVA, I.N.

Scattering of light in certain synthetic crystals. Kristallografiia
8 no.6:884-888 N-D'63. (MIRA 17:2)

1. Institut kristallografii AN SSSR.

MELANKHOLIN, N.M.

Absorption spectra of liquid crystals of thiazine dyes.
Opt. i spektr. 15 no.6:781-784 D '63. (MIRA 17:1)

MELANKHOLIN, N.M.

Pattern of growth and properties of liquid crystals of
thiazine dyes. Rost krist. 4:61-67 '64. (MIRA 17:8)

1 42396-65 EWT(1)/EMP(a)/EWT(m)/EMP(1)/T/EMP(t)/EMP(b)/EMA(c)/EEC(b).2 P1-4
 ACCESSION NR: AP5008485 IJP(c) JD/JG/GG/WH 8/0070/95/010/002/0214/0218 41

AUTHOR: Melankholin, N. M.; Martynova, N. G.

TITLE: On the nature of optical nonuniformities in ruby crystals 40
 B

SOURCE: Kristallografiya, v. 10, no. 2, 1965, 214-218

TOPIC TAGS: ruby optic material, crystal inhomogeneity, refractive index, interferometry, shadowgraph 21

ABSTRACT: A luminescent point source and Twyman interferometer were used to study nonuniformities in the refractive indices of ruby rods and plates. The studies were done in polarized light. Shadow patterns of the rods, made with the luminescent point source show two systems of bright bands perpendicular to and parallel with the optical axis of the crystal. These bands were found to correspond principally to mosaic blocks and partially to slip tracks. Shadow patterns were also made of growth layers, usually perpendicular to the axis of growth, but sometimes making less than a 90° angle. The gradual change in the refractive index throughout the specimen was studied with the Twyman interferometer. It was found that these changes are caused chiefly by irregularities in the distribution of chromium. 21

Card 1/2

L 42396-65

ACCESSION NR: AP5008465

When the chromium content in the specimens is increased, there is an increase in the radial gradient of the refractive index. The edges of mosaic blocks may be seen in interference patterns photographed in polarized light. These appear principally in patterns for the extraordinary wave. Jumps in the extraordinary refractive index as well as stresses at the edges of the blocks cause zigzags in the interference bands. These zigzags are sometimes as much as the width of a band, which means a jump in the refractive index of approximately $4 \cdot 10^{-6}$. Only in rare cases are the edges of the mosaic blocks visible in the interference pattern for the ordinary wave. In these cases, sometimes the edges perpendicular to the optical axis are seen. The conclusions drawn here apply to ruby crystals, however the method used is applicable to studies of nonuniformities in any crystals. Orig. art. has: 5 figures.

ASSOCIATION: Institut kristallografii AN SSSR (Institute of Crystallography, AN SSSR)

SUBMITTED: 19Jun64

ENCL: 00

SUB CODE: SS,OP

NO REF SOV: 003

OTHER: 000

Card 2/2

FUSHCHAROVSKIY, Yu.M.; MELANKHOLINA, Ye.N.

Characteristics of the tectonic structure of the Californian
Cenozoic folded zone. Trudy GIN no.89:55-119 '63.

(MIRA 18:6)

MELENKHOVINA, Ye.N.

Outline of the geology and tectonic development of the
Cenozoic tectonic zones in the western part of Oregon and
Washington (U.S.A.). Trudy G.N. no. 113:137-160 1964. (MIRA 18:9)

MELANKHOLINA, Ye.N.

Outline of the geology of Cretaceous sediments in the Maritime
Territory. Trudy GIN no.139:7-29 '65. (MIRA 18:9)

MILACINI, W.

"Ophthalmological Optics in Calculations", p. 68, (OPHTHALMOLOGICAL OPTICS, Vol. 42, 1945 (published 1952), Warsaw, Poland)

SC: Monthly List of East European Accessions (MEAL), 16, Vol. 1, No. 1, March 1955, Uncl.

MELANOWSKI, W.H.

Heredity in ophthalmology. Polski tygod.lek. 5 no.24:934-939 12
June 50. (CJML 20:5)

MELANOWSKI, W.H.

~~Basic problems of heredity in ophthalmology. Polski tygod. lek.~~
5 no.26:1021-1025; concl. 26 June 50. (CJML 20:5)

MELANOWSKI, W.H.

Allergy in eye diseases. Polski tygod. lek. 5 no.29-30:1089-1099
24 July 50. (CJML 20:5)

MELANOWSKI, W. H.

EXCERPTA MEDICA Sec.12 Vol.9/10 Ophthalmology Oct55

1606. MELANOWSKI W. H. Klin. Ocznej A. M., Warszawa. *Nowe dane z dziedziny neurookulistyki i ich znaczenie dla kliniki. Recent news from the field of neuro-ophthalmology and its significance for the clinic POST. OKULISTYKI 1954, 1 (19-34) Graphs 1 Illus. 15

The author calls more attention to the intracranial paths and optic centres. Presenting several of his own cases he points out the peculiarities and significance of this part of the visual apparatus. Finally he recommends careful examination of the visual field and colour sense especially while cooperating with the neurosurgeon.

Szmyt - Lódź

Melanowski, H.

EXCERPT A MEDICA Sec.12 Vol.9/7 Ophthalmology Jul 55

1043. MELANOWSKIEGO W. H. XXIV *Zjazd okulistów Polskich (14-16. X. 1954 r. Warszawa). XXIVth Congress of Polish Ophthalmologists, Warsaw 1954 PANSTW. ZAKŁ. WYDAWICTW. LEK. (Warszawa) 1954(3-144)

The first day was devoted to the problem of glaucoma presented by Prof. Sobanski. The author presented the results of personal experiments as well as the works of his clinical team under the title 'Experimental investigations on the outflow of the aqueous humour from the eyeball'. Dr. Segal discussed 'The pathogenesis of glaucoma'. The second day was devoted to the problem of 'Trauma of the eye' which was presented by Prof. Melanowski followed by Prof. Kapuściński who spoke about 'Sympathetic ophthalmia'. Prof. Krwawicz discussed 'The regeneration of the injured cornea and its histological picture'. The third day was devoted to various themes. Fifty-five papers were delivered and discussed. Szmyt - Łódź

EXCERPAT MEDICA Sec.12 Vol.11/10 Ophthalmology Oct57
MELANOWSKI W. H.

1575. MELANOWSKI W. H. Klin. Okulist. A. M., Warszawa. * Urazy narządu wzroku. Injuries of the eye POST. OKULIST. 1956, 3 (131-143) Illus. 18
Review based on personal experience and data from the literature. The question of the removal of non-magnetic foreign bodies from the eye as well as the treatment of thermal and chemical eye burns is amply discussed. The author holds the opinion that prophylactic methods are mostly to be recommended especially while working with some corrosive materials. Szmyt - Łódź

MELANOWSKI, Wladyslaw Henryk; KOBUSZEWSKA-FARYNA, Maria

Considerations on retrolental fibroplasia and its relation to hydrophthalmia. Klin. oczna 26 no.3:199-205 1956.

1. Z Kliniki Ocznej A.M. w Warszawie. Kierownik: prof. dr. W. H. Melanowski I z Zakladu Anatomii Patologicznej A.M. w Warszawie. Kierownik: prof. dr. L. Paskiewicz.
Adres autorow: Prof. W. H. Melanowski, W-wa, ul. Emilii Plater 35, m. 11. Doc. M. Kobuszevska-Faryna, W-wa, ul. Nobla 27, m. 6.

(HYDROPTHALMOS, complications,
retrolental fibroplasia (Pol))

(RETROLENTAL FIBROPLASIA, complications,
hydrophthalmos (Pol))

MELANOWSKI, Wladyslaw Henryk

Celebration of the 50th anniversary of Lagrange's sclerectomy
in Bordeaux. Klin. oczna 27 no.1:73-81 1957.

1. Warszawa, ul. Emilii Plater 35, m. 11.

(SCLERA, surg.

excis., 50th anniversary of Lagrange's operation (Pol))

MELANOWSKI, W.H.

Polish ophthalmology during the period of 15 years of Polish
People's Republic. Klin. oczna 29 no.3:229-231 '59.
(OPHTHALMOLOGY)